EIN European Journal of Neuroscience FENS

WILEY

Responsiveness to left-prefrontal tDCS varies according to arousal levels

Marco Esposito¹ ^(D) Carlo Miniussi³ ^(D)

| Clarissa Ferrari² | Claudia Fracassi¹ |
| Debora Brignani^{1,4} |

Revised: 3 December 2021

¹Neurophysiology Lab, IRCCS Istituto Centro San Giovanni di Dio Fatebenefratelli, Brescia, Italy

²Unit of Statistics, IRCCS Istituto Centro San Giovanni di Dio Fatebenefratelli, Brescia, Italy

³Center for Mind/Brain Sciences CIMeC, University of Trento, Rovereto, Italy

⁴Department of Clinical and Experimental Sciences, University of Brescia, Brescia, Italy

Correspondence

Debora Brignani and Claudia Fracassi, IRCCS Istituto Centro San Giovanni di Dio Fatebenefratelli, Via Pilastroni 4, 25125, Brescia, Italy. Email: debora.brignani@ cognitiveneuroscience.it; claudia.fracassi@cognitiveneuroscience.it

Funding information Italian Ministry of Health

Edited by: Gregor Thut

Abstract

Over the past two decades, the postulated modulatory effects of transcranial direct current stimulation (tDCS) on the human brain have been extensively investigated. However, recent concerns on reliability of tDCS effects have been raised, principally due to reduced replicability and to interindividual variability in response to tDCS. These inconsistencies are likely due to the interplay between the level of induced cortical excitability and unaccounted structural and state-dependent functional factors. On these grounds, we aimed at verifying whether the behavioural effects induced by a common tDCS montage (F3-rSOA) were influenced by the participants' arousal levels, as part of a broader mechanism of state-dependency. Pupillary dynamics were recorded during an auditory oddball task while applying either a sham or real tDCS. The tDCS effects were evaluated as a function of subjective and physiological arousal predictors (STAI-Y State scores and pre-stimulus pupil size, respectively). We showed that prefrontal tDCS hindered task learning effects on response speed such that performance improvement occurred during sham, but not real stimulation. Moreover, both subjective and physiological arousal predictors significantly explained performance during real tDCS, with interaction effects showing performance improvement only with moderate arousal levels; likewise, pupil response was affected by real tDCS according to the ongoing levels of arousal, with reduced dilation during higher arousal trials. These findings highlight the potential role of arousal in shaping the neuromodulatory outcome, thus emphasizing a more careful interpretation of null or negative results while also encouraging more individually tailored tDCS applications based on arousal levels, especially in clinical populations.

KEYWORDS

pupil, response variability, state dependency, tDCS

Abbreviations: LC, locus coeruleus; LMMs, linear mixed models; PD, pupil dilation; PrePD, pre-stimulus pupil diameter; RT, reaction times; STAI-Y, State–Trait Anxiety Inventory; tDCS, transcranial direct current stimulation.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made. © 2022 The Authors. *European Journal of Neuroscience* published by Federation of European Neuroscience Societies and John Wiley & Sons Ltd.

1 | INTRODUCTION

Founded on decades of experimentation, transcranial direct current stimulation (tDCS) is a research tool capable of interacting with the central nervous system that has been rediscovered at the beginning of this century (Priori, 2003). Beside its value for basic research (Antal et al., 2017), tDCS has raised great interest for real-world applications, like rehabilitative interventions for neurological and psychiatric diseases (Lefaucheur et al., 2017) and cognitive enhancement (or detraction) in both young and older adults (Ke et al., 2019; Nelson et al., 2016; Santarnecchi et al., 2015; Summers et al., 2016). However, our incomplete understanding on how the physiological effects link to a behavioural outcome has hindered the development of more effective and generalizable stimulation protocols (Bestmann et al., 2015). What raises most concern is the lack of replicability among studies and the interindividual variability in response to tDCS (Horvath et al., 2015a,b; Medina & Cason, 2017; Wiethoff et al., 2014). In addition to nonoptimal methodological practices, a complex interplay among brain anatomical differences (i.e., cortical surface topography, skull thickness, subcutaneous fat levels and cerebrospinal fluid density) and the level of neuromodulatory effects might be crucial in explaining the inconsistencies across studies. Furthermore, state-based factors, including the levels of activation prior and during stimulation, the performance rank, wakefulness, task priming or novelty, might all play a decisive role (Horvath et al., 2014; Krause & Kadosh, 2014; Li et al., 2015); likewise, trait-based differences that strictly depend on neurotransmitter systems, such as different degrees of impulsive and risky behaviour, are also key to understanding the potential negative effects of tDCS (Bell et al., 2020). As a result, it appears conceivable to interpret the final effects of tDCS as contingent on the level of network engagement (Fertonani & Miniussi, 2017; Miniussi et al., 2013). In line with this prediction, many experiments have demonstrated a clear effect of baseline levels of different mental capabilities on tDCS-induced response (Benwell et al., 2015; Berryhill & Jones, 2012; Hsu et al., 2016; Mauri et al., 2015; Sarkar et al., 2014). Most recently, variability in the context of prefrontal tDCS has been associated with GABA- and glutamatedriven levels of excitability of the targeted cortex (Filmer, Ehrhardt, Bollmann, et al., 2019).

Further concerns have been raised in the field of transcranial electric stimulation regarding the subcutaneous nerves stimulation that may contribute even unconsciously to the final behavioural outcome (Liu et al., 2018). The possibility of indirectly affecting brain circuits likely adds on the participant's initial state- or

trait-based levels that make these measurements even more important for a better interpretation of the end results.Importantly, the use of tDCS is known to have an impact on large-scale brain systems extending well beyond the area under the stimulating electrode (Antal et al., 2011; Sandrini et al., 2020; Wagner et al., 2014; Weber et al., 2014). This approach translates into a lack of focality that resembles the spread of the noradrenergic modulatory action exerted by the locus coeruleus (LC), which arguably subtends arousal functions. This midbrain system is widely recognized for its role in shaping behavioural performance of primates (Aston-Jones et al., 1999; Berridge & Waterhouse, 2003; Sara & Bouret, 2012). A large body of evidence also suggests that the exogenous direct currents and the endogenous modulatory action on target cells share the same central mechanism of neuronal gain control (Aston-Jones & Cohen, 2005; Lafon et al., 2017; Moxon et al., 2007; Servan-Schreiber et al., 1990). Therefore, an interrelation between the two stimulating activities seems reasonable to the extent that whenever the contrast between activated and inhibited units becomes sufficiently increased or decreased any further added neuromodulation can likely spoil the expected results. In this respect, the human LC endogenous action measured by pupil dilation during response inhibition processes was shown to depend on whether basal neural excitability was modulated or not prior to task completion (Adelhöfer et al., 2019). Furthermore, a widespread release of noradrenaline could selectively interact with the local glutamate-driven levels of excitability, which have been described as generating feedback loops in tasks with different degrees of stimulus priority (see the GANE model, Mather et al., 2016). To this regard, we have recently showed that existing information-priority advantage of high saliency stimuli was differently modulated when applying transcranial electric stimulation in young and elderly individuals, according to their arousal levels at baseline (Esposito et al., 2021).

Given the above considerations, it appears evident that great part of the tDCS behavioural variability may stem from the interdependency between the induced cortical excitability and the varying levels of arousal experienced by participants. The aim of this study was to verify whether the behavioural and physiological responses induced by tDCS were somehow dependent on the participants' arousal levels. We chose a broadly used tDCS montage in the neuromodulation literature (e.g., Dedoncker et al., 2016; Lefaucheur et al., 2017) that supposedly target the prefrontal cortex in basic cognitive protocols (e.g., Mulquiney et al., 2011; Penolazzi et al., 2013; Plewnia et al., 2013; Sandrini et al., 2014; Savic et al., 2017), as well as in clinical trials (e.g., Palm et al., 2012; Smith et al., 2015; Suemoto et al., 2014).

The tDCS was applied during an auditory oddball task aimed to probe cognitive performance as a function of arousal levels (Beatty, 1982) and purposefully designed to keep participants' attention on task over uncertain periods (i.e., variable instead of fixed inter stimulus interval) in a way that online tDCS effects would be more subjected to a higher volatility of sustained attention (Langner & Eickhoff, 2013; Unsworth et al., 2018).

We tracked pupillary changes as a proxy for the LC modulatory action (Costa & Rudebeck, 2016; Murphy et al., 2011). We used reaction times (RT) and pupil dilation (PD) as a measure of LC phasic response to the relevant stimuli (target) and pre-stimulus pupil diameter (PrePD) as a physiological marker of the LC tonic discharge activity. Subjective arousal levels were also evaluated by means of State–Trait Anxiety Inventory (STAI-Y), an assessment tool for both trait and state factors relating to anxiety, from which we used state-related scores (Spielberger, 2010).

2 | MATERIALS AND METHODS

2.1 | Participants

Eighteen right-handed healthy participants (mean [SD] age = 23.7 [3.8]; 10 females; mean [SD] STAI-Y trait score = 44.9 [3.9]) took part in the experiment. Participants had no history of neurological or psychiatric illness and had normal or corrected-to-normal visual acuity. Ethical approval was obtained by the Ethics Committee of the IRCCS Centro San Giovanni di Dio Fatebenefratelli, Brescia, Italy. All participants were given written informed consent.

2.2 | Experimental design and procedure

A single-blind within-subject design was implemented. The testing sessions were organized in two days separated by at least 48 h in order to exclude any tDCS carryover effects. In each session, participants completed the task twice: at *baseline* (T1) without any electrodes mounted on their scalp and subsequently either during *sham* or *real* stimulation (T2) (Figure 1b). We collected behavioural and pupil data for the whole task duration (\sim 18 min).

Participants were randomly assigned and counterbalanced across two session-orders of tDCS protocol, and they were kept blind to the ongoing experimental condition (i.e., sham or real). The same participant was tested at around the same hour to control for any arousal variation due to the daily metabolic cycle and circadian rhythms (Oken et al., 2006). Moreover, we controlled that all participants were balanced in terms of amount of sleep, caffeine, alcohol and nicotine consumption. Participants seated in a soundproof dark room at the distance of about 55 cm from a 17-in LCD monitor and with the only source of light provided by a grey fixation cross. The auditory oddball task was presented using E-Prime presentation software (Schneider et al., 2001) by means of two constant-loudness speakers (Figure 1a).

In every task condition, there was a fixed total number of trials (420) of which 20% included targets (84) and 80% standards stimuli (336). The interstimulus interval was set to a range of 2.1-2.9 s and both stimuli lasted for 70 ms. In so doing, we ensured enough time (\sim 8 s) for the phasic pupil dilation to return to baseline before overlapping to the next target trial, that is, by playing at least three non-target tones between a target and the next one (Gilzenrat et al., 2010; Murphy et al., 2011). Along with a short training session, participants were instructed to readily press a button with their right index finger whenever detecting a target tone and to keep their gaze on the fixation cross throughout the task. Speed of response and gaze fixation were emphasized before each task execution. At the end of each experimental session, participants were given a questionnaire to rate the perceived sensations or discomforts that influenced their performance (Fertonani et al., 2015).

2.3 | tDCS protocol

A battery-driven current stimulator (Brain-STIM, EMS, Bologna, Italy) was used to deliver 1-mA (0.028 mA/cm^2) direct current stimulation via two rubber electrodes (35 cm^2) which were inserted inside two saline-soaked sponges. In order to ensure a stable impedance level as well as keeping skin sensations at the minimum, conductive electro-gel was also applied. The electrodes montage consisted in placing the anode over the area F3 of the EEG 10-20 system and the return (cathode) electrode over the right supra-orbital area (Figure 1c). The duration of the stimulation consisted of about 17 min (1040 s) with 15 s of currents fade-in and fade-out. Configuration of the sham condition included 15 s of fade-in, 10 s of actual current delivery and 15 s of fadeout given at the beginning of the experiment only (Figure 1d).

Mindful that the mere sensory stimulation could confound the expected arousal effects, we run a pilot



Sham

15 s 10 s 15 s

FIGURE 1 Study design and task paradigm. (a) Example of a trial sequence. (b) Overview of the experimental timeline, showing two testing sessions each one with two task conditions: baseline and stimulation. (c) Simulation results for the applied tDCS montage and parameters using SimNIBS toolbox (Saturnino et al., 2019). The colours denote the strength of the electric fields (independently of current flow directionality) simulated in a default head model. (d) Schematic representation of the stimulation protocol

experiment to validate our blind-controlled tDCS protocol and to exclude unexpected technical concerns due to the use of eye-tracker and the simultaneous current delivery.

E 0.1

0.201

Ten right-handed healthy participants were recruited prior to the main study (mean age [SD] = 26.4 [1.7]; 6 females; mean [SD] STAI-Y trait score = 44.7 [2.3]). All participants underwent, in a counterbalanced order, real and sham tDCS at rest in the same day, interposed by a pause of 15 min. Each tDCS protocol lasted about 5 min, a duration whose excitability modulations have been proved to return to baseline within the following 5 min (Nitsche & Paulus, 2000, 2001). We showed that neither participants were subjectively aware of the type of stimulation (i.e., sensation questionnaire and oral report) nor did their rate of full or half-eye blinks, known to tap into the dopaminergic and fatigue-related neural pathways (Stern et al., 1994), differ between sham $(8.07\% \pm 5.8)$ and real stimulation $(7.76\% \pm 5.9)$ ($t_{(9)} = 0.47$; p = 0.64). Despite the limited sample size, we deemed this evidence supportive for our within-subject study design.

2.4 | Pupil signal recording and preprocessing

Pupil diameter was recorded with an EyeLink 1000 Plus system (SR Research, Osgood, ON, Canada) and at 500-Hz sampling rate with left-monocular and pupil-CR tracking mode. Pupil signal was processed offline. A shape-preserving piecewise cubic interpolation method was chosen to interpolate values ranging from 70 ms before blink onset to 300 after blink offset. Epoch segmentation and baseline correction were then carried out (respectively for -1 s to +2.5 s and -800 to +200 ms relative to target onset). All epochs with a peak pupil diameter exceeding ± 2 mm were rejected (Murphy et al., 2011). We extracted two variables of interest: (i) pupil dilation (PD), as the peak value of the maximum dilation after targets presentation, and (ii) pre-stimulus pupil diameter (PrePD), as the mean of 1 s data prior to tone presentation.

2.5 | Statistical analyses

As expected, the nature of our oddball task caused ceiling effects in the correct responses for all conditions (accuracy rate >98%). All trials including either a false alarm or a missed response were removed from subsequent analyses, as well as trials corresponding to RT faster than 150 ms or exceeding 1.96 standard deviations from the mean (number rejected trials: M = 3.14, SD = 1.39). All valid RT were then log-transformed to the base *e* in order to ensure a normal distribution of the data.

We considered only trials having no missing values at the two main outcomes RT and PD, resulting in 52 trials overall. Importantly, these data points were not collapsed across conditions: *Trial* was included in the analyses as an independent fixed factor, and thus affording a greater reliability and robustness of the findings.

In order to study the effect of tDCS on the behavioural and physiological responses, we performed two linear mixed models (LMM) on RT and on PD as dependent variables. Individual (subject-specific) variation was accounted for by considering Subjects as random effect. Fixed effects, repeated within subjects, were specified for Condition (2 levels, real and sham), Time (2 levels, T1 and T2) and Trial (52 levels), whereas Order (2 subgroups, sham-real and real-sham) was considered as a betweensubject fixed effect. In addition, the interaction Condition \times Time was assessed. Post hoc comparisons were adjusted with Sidak correction for multiple comparisons. The use of all repeated measures (including the 52 levels of Trial) at each time point allowed to considerably increase the power of the tests on the assessed effects that is, notoriously, affected by the sample size (see, e.g., the degree of freedom-reported in subscript parenthesesof the statistics of each effect reported in Section 3). The use of LMMs allows to manage and use all the data available at each time point in a repeated measure framework, and this, in turn, allows to increase the statistical power of the analysed effects. A post hoc power analysis was performed to provide proofs of reliability and robustness of our LMM results. In detail, once the LMMs were performed, the power of each fixed effect of LMMs was

analysed by *powerSim* function of the *simr* package of the R software (R core Team, 2020, http://www.R-project. org/), which provides both the power of the estimated effect and the corresponding 95% confidence interval.

The above LMM was subsequently adjusted for subjective arousal (measured by STAI-Y State score) and for physiological arousal (evaluated by PrePD) in order to assess their effects on tDCS-induced modulation. Akaike information criteria (AIC) was used to select the best fitted models (the lower AIC the better model) and the corresponding predictors.

3 | RESULTS

3.1 | Reaction times

The unadjusted linear mixed model on reaction times (RT) (AIC = -2545.34) revealed no significant effects of the Order $(F_{(1,16)} = 0.071, p = 0.794)$ and Trial $(F_{(51,126)} = 1.10, p = 0.322)$. A significant effect of *Time* $(F_{(1,2999)} = 4.53, p = 0.033)$ showed that performance significantly improved from T1 to T2 sessions, indicating an overall practice effect. In addition to a significant effect of *Condition* ($F_{(1,3040)} = 11.40$, p = 0.001), we found a significant *Condition* \times *Time* interaction effect $(F_{(1,3014)} = 10.96, p = 0.001)$, indicating a different trend for real and sham conditions. The post hoc comparison within the factor Time revealed a significant performance improvement during sham (p < 0.001), but not during *real* stimulation (p = 0.95). This finding suggests that real tDCS hindered the practice effect that was present in the sham condition.

Next, LMM adjusted for STAI-Y and PrePD were separately performed (see Table 1). We found an overall significant contribution of STAI-Y ($F_{(1,2886)} = 7.95$, p = 0.005) and more importantly a significant three-way interaction (Condition \times Time \times STAI-Y: $F_{(3,2108)} = 12.53$, p < 0.001), indicating that the subjective level of arousal affected the interaction Condition \times Time on RT in different way across time (T1 and T2) and conditions. Specifically, STAI-Y state scores were predictive of the performance variations across tDCS conditions: During sham session, a performance improvement (i.e., a decrease of RT in T2 stimulation) was observed for almost all the levels of arousal, and it visibly diminished at the higher end of the STAI-Y range. Differently, in the real tDCS condition, the improvement of RT reversed in a worsening pattern, with a more substantial performance decline (i.e., an increase of RT in T2 stimulation) as STAI-Y scores increased (see Figure 2 for a graphical display of the Condition \times Time \times STAI-Y significant effect).

TABLE 1 The structure and statistics of the models utilized for the dependent variable reaction time

behavioural performance (KT)						
Models	AIC	Fixed factors	F _(df)	<i>p</i> value		
(1) Unadjusted model	-2545	Trial	1.10(51,126)	0.322		
		Time ^a	$4.53_{(1,2999)}$	0.033		
		Order	0.07 _(1,15)	0.794		
		Condition ^a	11.40(1,3040)	0.001		
		$\mathit{Condition} imes \mathit{Time}^{\mathrm{a}}$	10.96(1,3014)	0.001		
(2) Adjusted for STAI-Y	-2536	STAI-Y ^a	$7.95_{(1,2886)}$	0.005		
		Trial	1.09(51,130)	0.334		
		Time ^a	24.83 _(1,2864)	< 0.001		
		Order	$0.11_{(1,16)}$	0.74		
		Condition	$1.74_{(1,2868)}$	0.187		
		$\textit{Condition} \times \textit{Time} \times \textit{STAI-Y}^{a}$	12.53 _(3,2108)	< 0.001		
(3) Adjusted for PrePD	-2519	PrePD	2.80(1,2462)	0.094		
		Trial	$1.02_{(51,124)}$	0.483		
		Time	$1.47_{(1,2734)}$	0.225		
		Order	$0.10_{(1,16)}$	0.745		
		Condition ^a	8.54 _(1,2903)	0.003		
		$\textit{Condition} \times \textit{Time} \times \textit{PrePD}^{a}$	5.40 _(3,1772)	0.001		

Behavioural performance (RT)

6

Note: All models include subjects as random factor. Significant fixed factors are displayed with an asterisk. ^aSignificant effect.

After adjusting for PrePD, the effect of Condition remained significant $(F_{(1,21903)} = 8.54, p = 0.003)$, but this was not the case for neither the effect of Time nor the physiological predictor ($F_{(1,2734)} = 1.47$, p = 0.225; $F_{(1,2462)} = 2.80, p = 0.094$, respectively). However, the significant three-way interaction effect (Con*dition* × *Time* × PrePD: $F_{(3,1772)} = 5.40$, p = 0.001) revealed that the participants' physiological level of arousal affected the RT in a different way across condition and time: Performance improvement across time was consistent in the sham condition, whereas RT became slower as ongoing physiological arousal increased (see Figure 3 for a graphical display of Condition \times Time \times PrePD significant effect).

By the post hoc power analysis, a power higher than 0.8 was found for all significant effects of the mixed models above reported, except for the effect of *Time* on reaction times ($F_{(1,2999)} = 4.53$, p = 0.033; power = 0.5 [95% CI: 23–65%]).

Overall, based on these data, a different trend emerged from the comparison of the investigated effects (i.e., RT change) between adjusted and unadjusted models (Figure 4). When using our adjusted models, the evident reduced variability in the direction of RT change corroborates the importance of not disregarding discrepancies rooted in interindividual differences, such as in physiological and subjective arousal, but rather include them as predictors along with individual random effects.

3.2 | Pupil dilation

In the unadjusted LMM on pupil dilation (PD), AIC = 857.37, all fixed effects were significant (*Condition*: $F_{(1,2069)} = 10.30$, p = 0.001; *Time*: $F_{(1,2102)} = 34.58$, p < 0.001; *Trial*: $F_{(51,121)} = 6.06$, p < 0.001) except for the factor *Order* ($F_{(1,15)} = 0.59$, p = 0.45) and the interaction between *Condition* and *Time* ($F_{(1,2080)} = 2.94$, p = 0.086). Importantly, pupil dilation decreased from *T1* (M = 0.36; SE = 0.021) to *T2* sessions (M = 0.31; SE = 0.021), indicating a general habituation of the phasic pupillary responses. However, no specific effect of tDCS on PD was revealed.

The adjustment for STAI-Y got slightly worse the model fitting (AIC = 872.86) (see Table 2), though with a significant effect of the predictor FIGURE 2 Reaction times by subjective arousal, condition and time. Average log-based RT of model fitted values is plotted as a function of STAI-Y scores (binned values), with results from session sham (top panel) and real (bottom panel) in the two time points T1 and T2. Each mean value is marked over the corresponding distribution of the data. Colours light green and orange represent the baseline (T1) and stimulation (T2) task, respectively. The figure aims to provide graphical evidence for the interaction term: Condition \times Time \times STAI-Y. For this purpose, the continuous variable STAI-Y was categorized in five binned values. In the session sham: Baseline RTs decreased as STAI-Y State scores increased (i.e., baseline RTs at ≤40 score were slower than RTs at the other scores, all ps < 0.001). No difference in RTs were found for T2 evaluation across the STAI-Y range, except for ≤ 40 versus ≥ 47 , p < 0.001. In the session real: Baseline RTs decreased as STAI-Y State scores increased (i.e., baseline RTs at ≤40 score were slower than RTs at the other scores, p < 0.001 for all). An increase in RTs was found for T2 evaluation across the STAI-Y range

6.1 RT log 6.05 6 5.95 T1>T2 T1>T2 T1>T2 T1>T2 T1<T2 5.9 p<0.001 p<0.001 p<0.001 p<0.001 p<0.001 5.85 ≤40 41-42 43-44 45-46 T1 baseline T2 stimulation Session REAL 6.15 6.1 RT log 6.05 6 5.95 T1>T2 T1<T2 T1<T2

European Journal of Neuroscience

FENS

Session SHAM

FINI

6.15

(i.e., T2 RTs at \leq 40 score were faster than RTs at other scores, all ps < 0.001, except for the 41-42 score)

T1>T2

p<0.001

≤40

5.9

5.85

 $(F_{(1,1744)} = 17.92, p < 0.001)$ and a significant interaction *Condition* × *Time* × STAI-Y ($F_{(3,1521)} = 3.01$, p = 0.029), suggesting that pupil dilation was also affected by tDCS according to the subjective levels of arousal.

Adjusting for PrePD strongly improved the model significant fitting (AIC = -658.59),with PrePD $(F_{(1,2548)} = 2337.89, p < 0.001)$ and interaction Con $dition \times Time \times PrePD$ effects $(F_{(3,1517)} = 12.58,$ p < 0.001). In detail, during the sham condition, a decrease in pupil dilation consistently occurred throughout the range of PrePD values, whereas during real tDCS, the pupil dilation progressively shifted towards a greater reduction during trials with larger PrePD (see Figure 5 for a graphical display of Condition \times Time \times PrePD significant effect).

By the post hoc power analysis, a power higher than 0.8 was found for all significant effects of the mixed models, except for effect the interaction Condition \times Time \times STAI-Y pupil dilation on $(F_{(3,1521)} = 3.01, p = 0.029; \text{ power } 0.6 [95\% \text{ CI: } 36-81]).$

DISCUSSION 4

p<0.001

41-42

p<0.001

43-44

STAI-Y State

p<0.001

45-46

In the present study, we addressed the question of whether variable effects of a single tDCS session could be dependent on the degree of arousal experienced before (i.e., subjective report of anxiety at baseline, STAI-Y) and during (i.e., physiological tonic LC activity, PrePD) the experiment. We found that subjective and physiological levels of arousal significantly accounted for the variation of reaction times across two experimental sessions. Real tDCS appeared to hinder the practice effect observed during the sham condition, with a trend becoming especially evident at higher levels of arousal. As for pupil dilation, its modulations matched with the behavioural results, with a more reduced pupillary response emerging during real tDCS as arousal levels increased.

These results shed light on one relevant factor, which may account for the paucity of consistency across tDCS effects: Arousal appears to be predictive of the

WILEY

≥47

T1<T2

p<0.001

≥47



ESPOSITO ET AL.

FIGURE 3 Reaction times by physiological arousal, condition and time. On each box, the interquartile range, the whiskers and the median of predicted log-based RT are represented for three linearly interspaced bins of pre-target pupil diameter, with results from session sham (top panel) and real (bottom panel) in the two time points T1 and T2. Colours light green and orange represent the baseline (T1) and stimulation (T2) task, respectively. The figure aims to provide graphical evidence for the interaction term: Condition \times Time \times PrePD. For this purpose, the continuous variable PrePD was categorized in three binned values. In the session sham: A decrease in RTs was found for the T1 evaluation across the PrePD bins (i.e., T1 RTs at first bin were slower than RTs at other bins, all ps < 0.001). A decrease in RTs was found for the T2 evaluation across the PrePD bins (i.e., T2 RTs at first bin were slower than RTs at other bins, all $p_{\rm S} < 0.05$). In the session *real*: for the T1 evaluation RTs increased only comparing the first bin to the third bin (p < 0.05). An increase in RTs was found for T2 evaluation across all the PrePD bins (i.e., T2 RTs at first bin were faster than RTs at other bins, all *ps* < 0.01)

modulations induced by tDCS on task performance. A number of studies, which reported a considerable interand intra-individual variability in response to tDCS protocols, investigated the impact of demographic characteristics (e.g., age and gender), cortical architecture variations or physiological measures specific to the targeted areas (e.g., levels of excitability of the primary motor cortex), yet without considering general measures of activation comparable to arousal (Filmer, Ehrhardt, Shaw, et al., 2019; Katz et al., 2017; López-Alonso et al., 2015; Ridding & Ziemann, 2010; van de Ruit & Grey, 2019; Wiethoff et al., 2014). Here, we collected ratings on the subjective level of anxiety (i.e., STAI-Y State) before each experimental session. Due to the close relation between the LC activity and the perceived anxiety (Eysenck, 1963; Mizuki et al., 1997; Robbins & Everitt, 1995), we used it as a fixed measure of arousal. Pre-target pupil diameter was instead used as a dynamic proxy of arousal, allowing us to track its ongoing fluctuations (Murphy et al., 2011; Unsworth et al., 2018; Van Den Brink et al., 2016). We confirmed that pupil dilation

values were negatively related with pre-target pupil diameter across all conditions, as frequently reported in the literature (De Gee et al., 2014; Hong et al., 2014; Murphy et al., 2011). Remarkably, despite the limited sample size, the large number of repeated measures properly used in a linear mixed model fashion allowed us to obtain an adequate power and thus robustness of our results.

When our measures of arousal were accounted for by statistical analyses, a clearer picture emerged, indicating that the effects induced by tDCS on the behavioural responses interacted with both subjective and physiological levels of arousal. Participants speeded up their responses when they completed the task for the second time in the sham session. This practice effect emerged somewhat independently of the ongoing physiological levels of arousal, although a more pronounced improvement appeared with lower levels of subjective arousal. During the application of real tDCS, however, performance ceased to improve more dramatically and, in fact, reversed with the exception of trials characterized by smaller pre-target pupil diameter and participants with a



FIGURE 4 Subject variability of reaction time change. Average log-based RT differences between the baseline (T1) and stimulation (T2) tasks are plotted on the vertical axis for each participant, separately for session sham (green bars) and real (orange bars). Each different bar plot is used to represent mean differences from the unadjusted (left panel) and the adjusted models using STAI-Y (middle panel) and PrePD (right panel) predictors. Negative and positive values on the horizontal axis indicate slower and faster performance, respectively

lower score at the STAI-Y questionnaire. A negative or null behavioural outcome of anodal tDCS is not uncommon in the literature and learning impairments have been reported in a host of different tDCS studies involving specific learning outcomes, such as unimproved working memory for recognition or implicit categorization, blocked consolidation of visual perception and inhibited motor learning (Ambrus et al., 2011; Berryhill et al., 2010; Bortoletto et al., 2015; Fertonani et al., 2011; Peters et al., 2013; Verhage et al., 2017). We chose response speed as behavioural measure, given that its intrinsic low sensitivity heavily relies on prior levels of fatigue and general activation (Welford, 1980). The interpretation of our behavioural results could be partially consistent with an inverted U-shape curve between task performance and arousal. According to this relationship, performance decline would occur when arousal levels are either too high or too low (Yerkes & Dodson, 1908). However, because none of the participants reported sleep deprivation or otherwise drowsiness-related conditions, we can assume that the lower values of our predictors effectively corresponded to moderate and not low levels of arousal. With this in

mind, the finding that facilitatory effects are principally associated with a moderate level of cortical excitation also seems to support the proposed cellular mechanism for a cortical excitation-inhibition balance (Krause et al., 2013). On these grounds, tDCS exogenous modulation would negatively impact on the normal cortical functioning whenever the levels of endogenous neural activity increase to the extent of a dysfunctional neuronal gain, with spontaneous task disengagement causing slower responses (Smallwood et al., 2004). A direct consequence of this mechanism would be the inhibition of task learning effects, unless the endogenous system is sufficiently inactive, as in low arousal trials. The latter scenario would provide an argument for when single session tDCS is found to improve task performance in the face of variable but otherwise moderate and wellbalanced arousal levels. Moreover, the understanding that an unbalanced combination of endogenous and exogenous excitability-increase events may lead to negative effects is also coherent with frameworks on brain activity-dependent plasticity and on signal-to-noise ratio mechanisms (Bortoletto et al., 2015; Fertonani & Miniussi, 2017).

WILEY EIN European Journal of Neuroscience FENS

TABLE 2 The structure and statistics of the models utilized for the dependent variable pupil dilation

Pupil dilatation (PD)						
Models	AIC	Fixed factors	F _(df)	p value		
(1) Unadjusted model	857	Trial ^a	6.06(51,121)	< 0.001		
		Time ^a	34.58(1,2102)	< 0.001		
		Order	0.59 _(1,15)	0.451		
		Condition ^a	10.30(1,2069)	0.001		
		Condition $ imes$ Time	2.94 _(1,2080)	0.086		
(2) Adjusted for STAI-Y	872	STAI-Y ^a	17.92 _(1,1744)	< 0.001		
		Trial ^a	6.13 _(51,118)	< 0.001		
		Time ^a	$3.98_{(1,2069)}$	0.046		
		Order	$1.44_{(1,16)}$	0.247		
		Condition	$0.06_{(1,2009)}$	0.807		
		$\textit{Condition} \times \textit{Time} \times \textit{STAI-Y}^a$	3.01 _(3,1521)	0.029		
(3) Adjusted for PrePD	-686	PrePD ^a	2337.89 _(1,2548)	< 0.001		
		Trial ^a	2.29 _(51,119)	< 0.001		
		Time ^a	$19.05_{(1,2683)}$	< 0.001		
		Order ^a	7.84 _(1,15)	0.013		
		Condition ^a	7.85 _(1,2158)	< 0.001		
		$Condition \times Time \times PrePD^a$	12.58(3,1517)	< 0.001		

Note: All models include subjects as random factor. Significant fixed factors are displayed with an asterisk. ^aSignificant effect.

The above interpretations are accompanied by results on pupil dilation, which represents a physiological response to relevant stimuli. When the ongoing levels of arousal were considered in the analyses, a specific effect of tDCS on pupil dilatation was revealed. An overall reduction of pupil dilation occurred when participants completed the task for the second time, consistently with a physiological habituation effect that paralleled the practice effect seen in the behavioural results (Rankin et al., 2009; Thompson & Spencer, 1966). In particular, pupil dilation evenly decreased for the entire range of arousal in the sham session, but different variations emerged during the application of real tDCS: As compared with the lower end of the arousal range, a more pronounced reduction in pupil dilation was observed in trials associated with higher physiological arousal. These, in fact, fell towards the same right end of the scale as the trials of unimproved response times during real tDCS. Therefore, habituation of a phasic response may not necessarily match the outcome direction of the behavioural improvement (Mackworth, 1968). Pupil dilations primarily reflect the timely increase of neural gain control, which translates into a system's responsivity amplification, and as such can be ascribed in the aforementioned inverted-U curve (Aston-Jones & Cohen, 2005; Hong et al., 2014; Murphy et al., 2011). The implication would

additive effect of an exogenous he that the neuromodulation would, on the one hand, contrast the natural habituation effect on pupil dilation occurring below the intermediate range of tonic arousal and, on the other hand, accentuate task disengagement at higher levels of tonic arousal, hence a greater reduction in phasic response. An analogous explanation was put forward in a recent tDCS work showing a reduction of pupil dilation-but no behavioural effects-during a Go-NoGo task, whereby it was argued that an offline tDCS enhancement of neuronal membrane potential could hinder or replace the endogenous gain control mechanisms of locus coeruleus (Adelhöfer et al., 2019): When participants were asked to withhold their responses, a correlation between the theta band inhibitory activity and pupil dilation was present during the sham stimulation, but not during real tDCS condition.

Furthermore, outside the tDCS literature, phasic pupillary responses were found to be reduced whenever participants' attention was not directed to the task, such as during episodes of mind wandering (Jubera-García et al., 2019; Mittner et al., 2014; Smallwood et al., 2011; Unsworth & Robison, 2016). Indeed, recent empirical and theoretical formulations of mind wandering have proposed that the locus coeruleus–norepinephrine system is tightly linked to different internally driven cognitive states, that



FIGURE 5 Pupil dilations explained by physiological arousal. Model fitted PD values are plotted against pre-target pupil diameter, with results from session sham (left panel) and real (right panel). Light green and orange best-fitting lines describe the trend of pupil dilation data points over pre-target pupil diameter respectively for the baseline (T1) and stimulation (T2) task. Dashed lines represent prediction functional bounds, that is, the uncertainty of predicting the fitted lines. The figure aims to provide a graphical evidence for the interaction term: Condition × Time × PrePD. In this case, the continuous variable PrePD needed not be categorized in bins because of the trial-by-trial correspondence with the PD values. Hence, we fitted a straight line as a way to better visualize the significant effect

is, on- and off-task states with various degrees of deliberate control (Christoff et al., 2016; Mittner et al., 2016). In this respect, the possibility of a direct and focally targeted tDCS modulation of mind wandering has been recently debated with uncertain conclusions (Chaieb et al., 2019). Based on this knowledge, it is not unlikely, although only hypothetical, that our tDCS effects would also be partly dependent on the arousal-mediated propensity of mind wandering activity during the task. Alternatively, mindfulness disposition, which is often considered to be in anticorrelation with mind-wandering, might lend itself to an analogous interpretation (Seli et al., 2016). Interestingly, it was recently found that tDCS decreased performance in a sustained attention task only in participants with high and not low dispositional mindfulness (Molina et al., 2020). This finding fits well with the idea of different internal states whose degree of intentionality and control always reflect the LC-driven transitions between explorative and exploitative mental operations (Mittner et al., 2016).

In summary, our data highlight a critical factor that could explain the negative or null effects of a common tDCS application. We are aware that these are preliminary results, and their interpretation may not directly generalize to all tDCS studies until further support is provided.

As arousal is considered a base layer for many highorder cognitive functions (Berridge & Waterhouse, 2003), our rationale behind the stimulation set-up did not include the targeting of a precise region of interest. Therefore, this approach differs from what is commonly used in several tDCS studies where the expectation is to define and/or rely upon topographical maps that support brain functions. The lack of an active control condition to exclude montage-specific effects may be considered a limitation of this study, and future research might find appropriate to investigate how and if different electrode arrangements and the involvement of specific functional areas can affect the interactive influence of arousal. On the one hand, a bold hypothesis would be that, regardless of the stimulation site, online tDCS affects the cortical interaction with endogenous modulations e.g., as measured by a behavioural index of phasic arousal (Luna et al., 2020). Hypothetically, a similar effect could be due to different cortical regions embedded in the same longrange specialized network that equally subserve a given cognitive task; alternatively, or in addition to this, we posit that the neural underpinning of arousal-for example, putative widespread noradrenergic projections in the brain-could potentially make the interaction with the induced cortical excitability irrespective of the spatial

11

FENS

ESPOSITO ET AL.

localization. On the other hand, though, the direction of the present results may have well depended on the specific interplay between tDCS montage, stimulation site and task. For example, it is possible that for most participants the current flowing through the anode and the cathode affected the orbitofrontal cortex, which is a key area in decision-making, and might have also played a crucial role in our task. The focus of future research should employ high-definition montages as well as carefully considering electric field intensity, focality and direction. Furthermore, the reciprocal interactions between the arousal system's neurotransmitters (e.g., noradrenaline and dopamine) should also be investigated whenever the stimulation site comprises parts of common projections (e.g., prefrontal regions, Briand et al., 2007).

5 | CONCLUSION

The large differences that we found in arousal levels stimulate reflection on what may mask the desired effects in the varied and still growing landscape of neurostimulation studies, which often fail to incorporate, but simply acknowledge, the crucial aspect of individual state-dependent variables (Lanina et al., 2018; Penton et al., 2018; Talsma et al., 2017; Willis et al., 2019). The importance of brain state is not a novel idea in the literature on non-invasive brain stimulation. The ongoing or basal levels of activation, included in the concept of "state-dependency," have been extensively reported to impact the effects of transcranial magnetic stimulation (TMS) (Silvanto et al., 2008) as well as tDCS (Antal et al., 2007). Considering the mechanisms of action of tDCS, which modulates excitability of neurons by hyperpolarizing or depolarizing their membrane potential (Liebetanz et al., 2002; Nitsche et al., 2003), tDCS effects might be even more sensitive to the arousal levels than TMS. In a similar vein, these considerations might be applicable to any kind of current stimulation modality.

Future tDCS studies might also consider useful to have both dynamic and fixed measures of arousal as an accurate way to monitor its impact on the final outcome. If successful, these achievements would be of great help also in assessing the degree of effectiveness with which tDCS protocols are being utilized to treat or ameliorate clinical conditions.

ACKNOWLEDGEMENT

This study was supported by the Italian Ministry of Health (Ricerca Corrente).

CONFLICT OF INTEREST

The authors declare no competing financial interests.

AUTHOR CONTRIBUTIONS

M.E. was responsible for the study concept and design, data acquisition and analyses and data interpretation and wrote the original draft. C.Fe. was responsible for the data analyses, data interpretation and writing, review and editing. C.Fr. was responsible for the study concept and design, data acquisition and data interpretation. C.M. was responsible for the data interpretation and writing, review and editing. D.B. was responsible for the study concept and design, data interpretation, writing, review and editing and supervision.

PEER REVIEW

The peer review history for this article is available at https://publons.com/publon/10.1111/ejn.15584.

DATA AVAILABILITY STATEMENT

The database containing the raw data used in this study is available at data.mendeley.com, https://doi.org/10. 17632/v9kt5v8p2b.1

ORCID

Marco Esposito https://orcid.org/0000-0001-6984-8569 Clarissa Ferrari https://orcid.org/0000-0002-4101-6872 Claudia Fracassi https://orcid.org/0000-0002-8249-7959

Carlo Miniussi D https://orcid.org/0000-0002-5436-4745 Debora Brignani D https://orcid.org/0000-0003-0848-5164

REFERENCES

- Adelhöfer, N., Mückschel, M., Teufert, B., Ziemssen, T., & Beste, C. (2019). Anodal tDCS affects neuromodulatory effects of the norepinephrine system on superior frontal theta activity during response inhibition. *Brain Structure and Function*, 224(3), 1291–1300. https://doi.org/10.1007/s00429-019-01839-3
- Ambrus, G. G., Zimmer, M., Kincses, Z. T., Harza, I., Kovács, G., Paulus, W., & Antal, A. (2011). The enhancement of cortical excitability over the DLPFC before and during training impairs categorization in the prototype distortion task. *Neuropsychologia*, 49(7), 1974–1980. https://doi.org/10.1016/j. neuropsychologia.2011.03.026
- Antal, A., Alekseichuk, I., Bikson, M., Brockmöller, J., Brunoni, A. R., Chen, R., ... Paulus, W. (2017). Low intensity transcranial electric stimulation: Safety, ethical, legal regulatory and application guidelines. *Clinical Neurophysiology*, 128(9), 1774–1809. https://doi.org/10.1016/j.clinph.2017.06.001
- Antal, A., Terney, D., Poreisz, C., & Paulus, W. (2007). Towards unravelling task-related modulations of neuroplastic changes induced in the human motor cortex. *European Journal of Neuroscience*, 26(9), 2687–2691. https://doi.org/10.1111/j.1460-9568.2007.05896.x

EIN European Journal of Neuroscience FENS

- Antal, A., Polania, R., Schmidt-Samoa, C., Dechent, P., & Paulus, W. (2011). Transcranial direct current stimulation over the primary motor cortex during fMRI. *NeuroImage*, 55(2), 590–596. https://doi.org/10.1016/j.neuroimage.2010. 11.085
- Aston-Jones, G., & Cohen, J. D. (2005). Adaptive gain and the role of the locus coeruleus-norepinephrine system in optimal performance. *Journal of Comparative Neurology*, 493(1), 99–110. https://doi.org/10.1002/cne.20723
- Aston-Jones, G., Rajkowski, J., & Cohen, J. (1999). Role of locus coeruleus in attention and behavioral flexibility. *Biological Psychiatry*, 46(9), 1309–1320. https://doi.org/10.1016/S0006-3223 (99)00140-7
- Beatty, J. (1982). Task-evoked pupillary responses, processing load, and the structure of processing resources. *Psychological Bulletin*, 91(2), 276–292. https://doi.org/10.1037/0033-2909.91.2.276
- Bell, S. B., Turner, B., Sawaki, L., & DeWall, N. (2020). When brain stimulation backfires: The effects of prefrontal cortex stimulation on impulsivity. *Social Cognitive and Affective Neuroscience*, nsaa049. https://doi.org/10.1093/scan/nsaa049
- Benwell, C. S. Y., Learmonth, G., Miniussi, C., Harvey, M., & Thut, G. (2015). Non-linear effects of transcranial direct current stimulation as a function of individual baseline performance: Evidence from biparietal tDCS influence on lateralized attention bias. *Cortex*, 69, 152–165. https://doi.org/10.1016/j. cortex.2015.05.007
- Berridge, C. W., & Waterhouse, B. D. (2003). The locus coeruleusnoradrenergic system: Modulation of behavioral state and state-dependent cognitive processes. *Brain Research Reviews*, 42(1), 33–84. https://doi.org/10.1016/S0165-0173(03)00143-7
- Berryhill, M. E., & Jones, K. T. (2012). tDCS selectively improves working memory in older adults with more education. *Neuroscience Letters*, 521(2), 148–151. https://doi.org/10.1016/j. neulet.2012.05.074
- Berryhill, M. E., Wencil, E. B., Branch Coslett, H., & Olson, I. R. (2010). A selective working memory impairment after transcranial direct current stimulation to the right parietal lobe. *Neuroscience Letters*, 479(3), 312–316. https://doi.org/10.1016/j. neulet.2010.05.087
- Bestmann, S., de Berker, A. O., & Bonaiuto, J. (2015). Understanding the behavioural consequences of noninvasive brain stimulation. *Trends in Cognitive Sciences*, 19(1), 13–20. https://doi. org/10.1016/j.tics.2014.10.003
- Bortoletto, M., Pellicciari, M. C., Rodella, C., & Miniussi, C. (2015). The interaction with task-induced activity is more important than polarization: A tDCS study. *Brain Stimulation*, 8(2), 269– 276. https://doi.org/10.1016/j.brs.2014.11.006
- Briand, L. A., Gritton, H., Howe, W. M., Young, D. A., & Sarter, M. (2007). Modulators in concert for cognition: Modulator interactions in the prefrontal cortex. *Progress in Neurobiology*, *83*(2), 69–91. https://doi.org/10.1016/j.pneurobio.2007.06.007
- Chaieb, L., Antal, A., Derner, M., Leszczyński, M., & Fell, J. (2019). New perspectives for the modulation of mind-wandering using transcranial electric brain stimulation. *Neuroscience*, 409, 69– 80. https://doi.org/10.1016/j.neuroscience.2019.04.032
- Christoff, K., Irving, Z. C., Fox, K. C. R., Spreng, R. N., & Andrews-Hanna, J. R. (2016). Mind-wandering as spontaneous thought: A dynamic framework. *Nature Reviews Neuroscience*, 17(11), 718–731. https://doi.org/10.1038/nrn.2016.113

- Costa, V. D., & Rudebeck, P. H. (2016). More than meets the eye: The relationship between pupil size and locus coeruleus activity. *Neuron*, *89*(1), 8–10. https://doi.org/10.1016/j.neuron.2015. 12.031
- De Gee, J. W., Knapen, T., & Donner, T. H. (2014). Decision-related pupil dilation reflects upcoming choice and individual bias. Proceedings of the National Academy of Sciences of the United States of America, 111(5), E618–E625. https://doi.org/ 10.1073/pnas.1317557111
- Dedoncker, J., Brunoni, A. R., Baeken, C., & Vanderhasselt, M. A. (2016). A systematic review and meta-analysis of the effects of transcranial direct current stimulation (tDCS) over the dorsolateral prefrontal cortex in healthy and neuropsychiatric samples: Influence of stimulation parameters. *Brain Stimulation*, 9(4), 501–517. https://doi.org/10.1016/j.brs.2016. 04.006
- Esposito, M., Mauri, P., Panizza, L., Mazza, V., Miniussi, C., & Brignani, D. (2021). Baseline levels of alertness influence tES effects along different age-related directions. *Neuropsychologia*, *160*, 107966. https://doi.org/10.1016/j.neuropsychologia.2021. 107966
- Eysenck, H. J. (1963). Biological basis of personality. *Nature*, 199(4898), 1031–1034. https://doi.org/10.1038/1991031a0
- Fertonani, A., Pirulli, C., & Miniussi, C. (2011). Random noise stimulation improves neuroplasticity in perceptual learning. *Journal of Neuroscience*, 31(43), 15416–15423. https://doi.org/10. 1523/JNEUROSCI.2002-11.2011
- Fertonani, A., Ferrari, C., & Miniussi, C. (2015). What do you feel if I apply transcranial electric stimulation? Safety, sensations and secondary induced effects. *Clinical Neurophysiology*, *126*(11), 2181–2188. https://doi.org/10.1016/j.clinph.2015. 03.015
- Fertonani, A., & Miniussi, C. (2017). Transcranial electrical stimulation: What we know and do not know about mechanisms. *The Neuroscientist*, 23(2), 109–123. https://doi.org/10.1177/ 1073858416631966
- Filmer, H. L., Ehrhardt, S. E., Bollmann, S., Mattingley, J. B., & Dux, P. E. (2019). Accounting for individual differences in the response to tDCS with baseline levels of neurochemical excitability. *Cortex*, 115, 324–334. https://doi.org/10.1016/j.cortex. 2019.02.012
- Filmer, H. L., Ehrhardt, S. E., Shaw, T. B., Mattingley, J. B., & Dux, P. E. (2019). The efficacy of transcranial direct current stimulation to prefrontal areas is related to underlying cortical morphology. *NeuroImage*, 196, 41–48. https://doi.org/10.1016/ j.neuroimage.2019.04.026
- Gilzenrat, M. S., Nieuwenhuis, S., Jepma, M., & Cohen, J. D. (2010). Pupil diameter tracks changes in control state predicted by the adaptive gain theory of locus coeruleus function. *Cognitive, Affective, & Behavioral Neuroscience, 10*(2), 252–269. https:// doi.org/10.3758/CABN.10.2.252
- Hong, L., Walz, J. M., & Sajda, P. (2014). Your eyes give you away: Prestimulus changes in pupil diameter correlate with poststimulus task-related EEG dynamics. *PLoS ONE*, 9(3), e91321. https://doi.org/10.1371/journal.pone.0091321
- Horvath, J. C., Carter, O., & Forte, J. D. (2014). Transcranial direct current stimulation: Five important issues we aren't discussing (but probably should be). *Frontiers in Systems Neuroscience*, 8, 2. https://doi.org/10.3389/fnsys.2014.00002

WILEY

WILEY EIN European Journal of Neuroscience FENS

- Horvath, J. C., Forte, J. D., & Carter, O. (2015a). Evidence that transcranial direct current stimulation (tDCS) generates littleto-no reliable neurophysiologic effect beyond MEP amplitude modulation in healthy human subjects: A systematic review. *Neuropsychologia*, 66, 213–236. https://doi.org/10.1016/j. neuropsychologia.2014.11.021
- Horvath, J. C., Forte, J. D., & Carter, O. (2015b). Quantitative review finds no evidence of cognitive effects in healthy populations from single-session transcranial direct current stimulation (tDCS). *Brain Stimulation*, 8(3), 535–550. https:// doi.org/10.1016/j.brs.2015.01.400
- Hsu, T.-Y., Juan, C.-H., & Tseng, P. (2016). Individual differences and state-dependent responses in transcranial direct current stimulation. *Frontiers in Human Neuroscience*, 10, 643. https:// doi.org/10.3389/fnhum.2016.00643
- Jubera-García, E., Gevers, W., & Van Opstal, F. (2019). Influence of content and intensity of thought on behavioral and pupil changes during active mind-wandering, off-focus, and on-task states. *Attention, Perception, and Psychophysics*, 1–11, 1125– 1135. https://doi.org/10.3758/s13414-019-01865-7
- Katz, B., Au, J., Buschkuehl, M., Abagis, T., Zabel, C., Jaeggi, S. M., & Jonides, J. (2017). Individual differences and long-term consequences of tDCS-augmented cognitive training. *Journal of Cognitive Neuroscience*, 29(9), 1498–1508. https://doi.org/10.1162/jocn_a_01115
- Ke, Y., Wang, N., Du, J., Kong, L., Liu, S., Xu, M., An, X., & Ming, D. (2019). The effects of transcranial direct current stimulation (tDCS) on working memory training in healthy young adults. *Frontiers in Human Neuroscience*, 13, 19. https://doi. org/10.3389/fnhum.2019.00019
- Krause, B., & Kadosh, R. C. (2014). Not all brains are created equal: The relevance of individual differences in responsiveness to transcranial electrical stimulation. *Frontiers in Systems Neuroscience*, 8, 25. https://doi.org/10.3389/fnsys.2014.00025
- Krause, B., Márquez-Ruiz, J., & Cohen Kadosh, R. (2013). The effect of transcranial direct current stimulation: A role for cortical excitation/inhibition balance? *Frontiers in Human Neuroscience*, 7, 602. https://doi.org/10.3389/fnhum.2013.00602
- Lafon, B., Rahman, A., Bikson, M., & Parra, L. C. (2017). Direct current stimulation alters neuronal input/output function. *Brain Stimulation*, 10(1), 36–45. https://doi.org/10.1016/j.brs.2016. 08.014
- Langner, R., & Eickhoff, S. B. (2013). Sustaining attention to simple tasks: A meta-analytic review of the neural mechanisms of vigilant attention. *Psychological Bulletin*, 139(4), 870–900. https:// doi.org/10.1037/a0030694
- Lanina, A. A., Feurra, M., & Gorbunova, E. S. (2018). No effect of the right posterior parietal cortex tDCS in dual-target visual search. *Frontiers in Psychology*, 9, 2112. https://doi.org/10. 3389/fpsyg.2018.02112
- Lefaucheur, J. P., Antal, A., Ayache, S. S., Benninger, D. H., Brunelin, J., Cogiamanian, F., ... Paulus, W. (2017). Evidencebased guidelines on the therapeutic use of transcranial direct current stimulation (tDCS). *Clinical Neurophysiology*, *128*(1), 56–92. https://doi.org/10.1016/j.clinph.2016.10.087
- Li, L. M., Uehara, K., & Hanakawa, T. (2015). The contribution of interindividual factors to variability of response in transcranial direct current stimulation studies. *Frontiers in Cellular Neuroscience*, 9, 181. https://doi.org/10.3389/fncel.2015.00181

- Liebetanz, D., Nitsche, M., Tergau, F., & Paulus, W. (2002). Pharmacological approach to the mechanisms of transcranial DCstimulation-induced after-effects of human motor cortex excitability. *Brain*, 125(10), 2238–2247. https://doi.org/10.1093/ brain/awf238
- Liu, A., Vöröslakos, M., Kronberg, G., Henin, S., Krause, M. R., Huang, Y., ... Buzsáki, G. (2018). Immediate neurophysiological effects of transcranial electrical stimulation. *Nature Communications*, 9(1), 1–12. https://doi.org/10.1038/s41467-018-07233-7
- López-Alonso, V., Fernández-del-Olmo, M., Costantini, A., Gonzalez-Henriquez, J. J., & Cheeran, B. (2015). Intraindividual variability in the response to anodal transcranial direct current stimulation. *Clinical Neurophysiology*, *126*(12), 2342–2347. https://doi.org/10.1016/j.clinph.2015.03.022
- Luna, F. G., Román-Caballero, R., Barttfeld, P., Lupiáñez, J., & Martín-Arévalo, E. (2020). A high-definition tDCS and EEG study on attention and vigilance: Brain stimulation mitigates the executive but not the arousal vigilance decrement. *Neuropsychologia*, 142, 107447. https://doi.org/10.1016/j. neuropsychologia.2020.107447
- Mackworth, J. F. (1968). Vigilance, arousal, and habituation. Psychological Review, 75(4), 308–322. https://doi.org/10.1037/ h0025896
- Mather, M., Clewett, D., Sakaki, M., & Harley, C. W. (2016). Norepinephrine ignites local hotspots of neuronal excitation: How arousal amplifies selectivity in perception and memory. *Behavioral and Brain Sciences*, 39(2016), e200. https://doi.org/10. 1017/S0140525X15000667
- Mauri, P., Miniussi, C., Balconi, M., & Brignani, D. (2015). Bursts of transcranial electrical stimulation increase arousal in a continuous performance test. *Neuropsychologia*, 74, 127–136. https:// doi.org/10.1016/j.neuropsychologia.2015.03.006
- Medina, J., & Cason, S. (2017). No evidential value in samples of transcranial direct current stimulation (tDCS) studies of cognition and working memory in healthy populations. *Cortex*, 94, 131–141. https://doi.org/10.1016/j.cortex.2017.06.021
- Miniussi, C., Harris, J. A., & Ruzzoli, M. (2013). Modelling noninvasive brain stimulation in cognitive neuroscience. *Neurosci*ence and Biobehavioral Reviews, 37(8), 1702–1712. https://doi. org/10.1016/j.neubiorev.2013.06.014
- Mittner, M., Boekel, W., Tucker, A. M., Turner, B. M., Heathcote, A., & Forstmann, B. U. (2014). When the brain takes a break: A model-based analysis of mind wandering. *Journal of Neuroscience*, 34(49), 16286–16295. https://doi.org/ 10.1523/JNEUROSCI.2062-14.2014
- Mittner, M., Hawkins, G. E., Boekel, W., & Forstmann, B. U. (2016). A neural model of mind wandering. *Trends in Cognitive Sciences*, 20(8), 570–578. https://doi.org/10.1016/j.tics.2016. 06.004
- Mizuki, Y., Suetsugi, M., Ushijima, I., & Yamada, M. (1997). Differential effects of dopaminergic drugs on anxiety and arousal in healthy volunteers with high and low anxiety. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 21(4), 573–590. https://doi.org/10.1016/S0278-5846 (97)00033-X
- Molina, E., Fernandez-Prieto, I., Andres, P., & Parmentier, F. (2020). Opposite effects of transcranial direct current stimulation (tDCS) on sustained attention in individuals with high

EIN European Journal of Neuroscience FENS

and low levels of dispositional mindfulness. https://doi.org/10. 31234/osf.io/8ynca

- Moxon, K. A., Devilbiss, D. M., Chapin, J. K., & Waterhouse, B. D. (2007). Influence of norepinephrine on somatosensory neuronal responses in the rat thalamus: A combined modeling and in vivo multi-channel, multi-neuron recording study. *Brain Research*, 1147(1), 105–123. https://doi.org/10.1016/j.brainres. 2007.02.006
- Mulquiney, P. G., Hoy, K. E., Daskalakis, Z. J., & Fitzgerald, P. B. (2011). Improving working memory: Exploring the effect of transcranial random noise stimulation and transcranial direct current stimulation on the dorsolateral prefrontal cortex. *Clinical Neurophysiology*, *122*(12), 2384–2389. https://doi.org/10. 1016/j.clinph.2011.05.009
- Murphy, P. R., Robertson, I. H., Balsters, J. H., & O'connell, R. G. (2011). Pupillometry and P3 index the locus coeruleusnoradrenergic arousal function in humans. *Psychophysiology*, 48(11), 1532–1543. https://doi.org/10.1111/j.1469-8986.2011. 01226.x
- Nelson, J., McKinley, R. A., Phillips, C., McIntire, L., Goodyear, C., Kreiner, A., & Monforton, L. (2016). The effects of transcranial direct current stimulation (tDCS) on multitasking throughput capacity. *Frontiers in Human Neuroscience*, 10, 589. https:// doi.org/10.3389/fnhum.2016.00589
- Nitsche, M. A., Fricke, K., Henschke, U., Schlitterlau, A., Liebetanz, D., Lang, N., Henning, S., Tergau, F., & Paulus, W. (2003). Pharmacological modulation of cortical excitability shifts induced by transcranial direct current stimulation in humans. *Journal of Physiology*, 553(1), 293–301. https://doi. org/10.1113/jphysiol.2003.049916
- Nitsche, M. A., & Paulus, W. (2000). Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation. *Journal of Physiology*, 527(3), 633–639. https://doi. org/10.1111/j.1469-7793.2000.t01-1-00633.x
- Nitsche, M. A., & Paulus, W. (2001). Sustained excitability elevations induced by transcranial DC motor cortex stimulation in humans. *Neurology*, 57(10), 1899–1901. https://doi.org/10. 1212/WNL.57.10.1899
- Oken, B. S., Salinsky, M. C., & Elsas, S. M. (2006). Vigilance, alertness, or sustained attention: physiological basis and measurement. *Clinical Neurophysiology*, *117*(9), 1885–1901. https://doi. org/10.1016/j.clinph.2006.01.017
- Palm, U., Schiller, C., Fintescu, Z., Obermeier, M., Keeser, D., Reisinger, E., Pogarell, O., Nitsche, M. A., Möller, H. J., & Padberg, F. (2012). Transcranial direct current stimulation in treatment resistant depression: A randomized double-blind, placebo-controlled study. *Brain Stimulation*, 5(3), 242–251. https://doi.org/10.1016/j.brs.2011.08.005
- Penolazzi, B., Pastore, M., & Mondini, S. (2013). Electrode montage dependent effects of transcranial direct current stimulation on semantic fluency. *Behavioural Brain Research*, 248, 129–135. https://doi.org/10.1016/j.bbr.2013.04.007
- Penton, T., Bate, S., Dalrymple, K. A., Reed, T., Kelly, M., Godovich, S., Tamm, M., Duchaine, B., & Banissy, M. J. (2018). Using high frequency transcranial random noise stimulation to modulate face memory performance in younger and older adults: Lessons learnt from mixed findings. *Frontiers in Neuroscience*, 12, 863. https://doi.org/10.3389/fnins.2018.00863

- Peters, M. A. K., Thompson, B., Merabet, L. B., Wu, A. D., & Shams, L. (2013). Anodal tDCS to V1 blocks visual perceptual learning consolidation. *Neuropsychologia*, 51(7), 1234–1239. https://doi.org/10.1016/j.neuropsychologia.2013.03.013
- Plewnia, C., Zwissler, B., Längst, I., Maurer, B., Giel, K., & Krüger, R. (2013). Effects of transcranial direct current stimulation (tDCS) on executive functions: Influence of COMT Val/-Met polymorphism. *Cortex*, 49(7), 1801–1807. https://doi.org/ 10.1016/j.cortex.2012.11.002
- Priori, A. (2003). Brain polarization in humans: A reappraisal of an old tool for prolonged non-invasive modulation of brain excitability. *Clinical Neurophysiology*, 114(4), 589–595. https://doi. org/10.1016/S1388-2457(02)00437-6
- Rankin, C. H., Abrams, T., Barry, R. J., Bhatnagar, S., Clayton, D. F., Colombo, J., Coppola, G., Geyer, M. A., Glanzman, D. L., Marsland, S., McSweeney, F. K., Wilson, D. A., Wu, C.-F., & Thompson, R. F. (2009). Habituation revisited: An updated and revised description of the behavioral characteristics of habituation. *Neurobiology of Learning and Memory*, 92(2), 135–138. https://doi.org/10.1016/ j.nlm.2008.09.012
- Ridding, M. C., & Ziemann, U. (2010). Determinants of the induction of cortical plasticity by non-invasive brain stimulation in healthy subjects. *Journal of Physiology*, 588(13), 2291–2304. https://doi.org/10.1113/jphysiol.2010.190314
- Robbins, T., & Everitt, B. (1995). Central norepinephrine neurons and behavior. *Psychopharmacology: The Fourth Generation of Progress*, 363–372.
- Sandrini, M., Brambilla, M., Manenti, R., Rosini, S., Cohen, L. G., & Cotelli, M. (2014). Noninvasive stimulation of prefrontal cortex strengthens existing episodic memories and reduces forgetting in the elderly. *Frontiers in Aging Neuroscience*, 6, 289. https:// doi.org/10.3389/fnagi.2014.00289
- Sandrini, M., Xu, B., Volochayev, R., Awosika, O., Wang, W. T., Butman, J. A., & Cohen, L. G. (2020). Transcranial direct current stimulation facilitates response inhibition through dynamic modulation of the fronto-basal ganglia network. *Brain Stimulation*, 13(1), 96–104. https://doi.org/10.1016/j.brs. 2019.08.004
- Santarnecchi, E., Brem, A.-K., Levenbaum, E., Thompson, T., Kadosh, R. C., & Pascual-Leone, A. (2015). Enhancing cognition using transcranial electrical stimulation. *Current Opinion in Behavioral Sciences*, 4, 171–178. https://doi.org/10.1016/j. cobeha.2015.06.003
- Sara, S. J., & Bouret, S. (2012). Orienting and reorienting: The locus coeruleus mediates cognition through arousal. *Neuron*, 76(1), 130–141. https://doi.org/10.1016/j.neuron.2012.09.011
- Sarkar, A., Dowker, A., & Kadosh, R. C. (2014). Cognitive enhancement or cognitive cost: Trait-specific outcomes of brain stimulation in the case of mathematics anxiety. *Journal of Neuroscience*, 34(50), 16605–16610. https://doi.org/10.1523/ JNEUROSCI.3129-14.2014
- Saturnino, G. B., Puonti, O., Nielsen, J. D., Antonenko, D., Madsen, K. H., & Thielscher, A. (2019). SimNIBS 2.1: A Comprehensive Pipeline for Individualized Electric Field Modelling for Transcranial Brain Stimulation. *In: Makarov S., Horner M., Noetscher G.* (eds). Springer, Cham: Brain and Human Body Modeling. https://doi.org/10.1007/978-3-030-21293-3_1

_WILEY

WILEY EIN European Journal of Neuroscience FENS

- Savic, B., Cazzoli, D., Müri, R., & Meier, B. (2017). No effects of transcranial DLPFC stimulation on implicit task sequence learning and consolidation. *Scientific Reports*, 7(1), 1–10. https://doi.org/10.1038/s41598-017-10128-0
- Schneider, W., Eschman, A., & Zuccolotto, A. (2001). E-Prime user's guide. Psychology Tools.
- Seli, P., Risko, E. F., Smilek, D., & Schacter, D. L. (2016). Mind-wandering with and without intention. *Trends in Cognitive Sciences*, 20(8), 605–617. https://doi.org/10.1016/j.tics. 2016.05.010
- Servan-Schreiber, D., Printz, H., & Cohen, J. D. (1990). A network model of catecholamiine effects: Gain, signal-to-noise ratio, and behavior. *Science*, 249(4971), 892–895. https://doi.org/10. 1126/science.2392679
- Silvanto, J., Muggleton, N., & Walsh, V. (2008). State-dependency in brain stimulation studies of perception and cognition. *Trends in Cognitive Sciences*, 12(12), 447–454. https://doi.org/ 10.1016/j.tics.2008.09.004
- Smallwood, J., Brown, K. S., Tipper, C., Giesbrecht, B., Franklin, M. S., Mrazek, M. D., Carlson, J. M., & Schooler, J. W. (2011). Pupillometric evidence for the decoupling of attention from perceptual input during offline thought. *PLoS ONE*, 6(3), e18298. https://doi.org/10.1371/ journal.pone.0018298
- Smallwood, J., Davies, J. B., Heim, D., Finnigan, F., Sudberry, M., O'Connor, R., & Obonsawin, M. (2004). Subjective experience and the attentional lapse: Task engagement and disengagement during sustained attention. *Consciousness and Cognition*, 13(4), 657–690. https://doi.org/10.1016/j.concog.2004.06.003
- Smith, R. C., Boules, S., Mattiuz, S., Youssef, M., Tobe, R. H., Sershen, H., Lajtha, A., Nolan, K., Amiaz, R., & Davis, J. M. (2015). Effects of transcranial direct current stimulation (tDCS) on cognition, symptoms, and smoking in schizophrenia: A randomized controlled study. *Schizophrenia Research*, *168*(1–2), 260–266. https://doi.org/10.1016/j.schres.2015.06.011
- Spielberger, D. C. (2010). State-Trait Anxiety Inventory. In *The Corsini encyclopedia of psychology* (pp. 1–1). John Wiley & Sons, Inc.. https://doi.org/10.1002/9780470479216.corpsy0943
- Stern, J. A., Louis, S., & Boyer, D. (1994). Blink rate: A possible measure of fatigue. *Human Factors*, 36(2), 285–297. https:// doi.org/10.1177/001872089403600209
- Suemoto, C. K., Apolinario, D., Nakamura-Palacios, E. M., Lopes, L., Paraizo Leite, R. E., Sales, M. C., Nitrini, R., Brucki, S. M., Morillo, L. S., Magaldi, R. M., & Fregni, F. (2014). Effects of a non-focal plasticity protocol on apathy in moderate Alzheimer's disease: A randomized, double-blind, sham-controlled trial. *Brain Stimulation*, 7(2), 308–313. https://doi.org/10.1016/j.brs.2013.10.003
- Summers, J. J., Kang, N., & Cauraugh, J. H. (2016). Does transcranial direct current stimulation enhance cognitive and motor functions in the ageing brain? A systematic review and meta- analysis. Ageing Research Reviews, 25, 42–54. https://doi. org/10.1016/j.arr.2015.11.004
- Talsma, L. J., Kroese, H. A., & Slagter, H. A. (2017). Boosting cognition: Effects of multiple-session transcranial direct current stimulation on working memory. *Journal of Cognitive Neuroscience*, 29(4), 755–768. https://doi.org/10.1162/jocn_a_01077
- Thompson, R. F., & Spencer, W. A. (1966). Habituation: A model phenomenon for the study of neuronal substrates of behavior.

Psychological Review, 73(1), 16-43. https://doi.org/10.1037/h0022681

- Unsworth, N., & Robison, M. K. (2016). Pupillary correlates of lapses of sustained attention. Cognitive, Affective, & Behavioral Neuroscience, 16(4), 601–615. https://doi.org/10.3758/s13415-016-0417-4
- Unsworth, N., Robison, M. K., & Miller, A. L. (2018). Pupillary correlates of fluctuations in sustained attention. *Journal of Cognitive Neuroscience*, 30(9), 1241–1253. https://doi.org/10.1162/ jocn_a_01251
- Van de Ruit, M., & Grey, M. J. (2019). Interindividual variability in use-dependent plasticity following visuomotor learning: The effect of handedness and muscle trained. *Journal of Motor Behavior*, 51(2), 171–184. https://doi.org/10.1080/00222895. 2018.1446125
- Van Den Brink, R. L., Murphy, P. R., & Nieuwenhuis, S. (2016). Pupil diameter tracks lapses of attention. *PLoS ONE*, 11(10), 1–16. https://doi.org/10.1371/journal.pone.0165274
- Verhage, M. C., Avila, E. O., Frens, M. A., Donchin, O., & van der Geest, J. N. (2017). Cerebellar tDCS does not enhance performance in an implicit categorization learning task. *Frontiers in Psychology*, 8, 476. https://doi.org/10.3389/fpsyg.2017.00476
- Wagner, S., Rampersad, S. M., Aydin, Ü., Vorwerk, J., Oostendorp, T. F., Neuling, T., Herrmann, C. S., Stegeman, D. F., & Wolters, C. H. (2014). Investigation of tDCS volume conduction effects in a highly realistic head model. *Journal of Neural Engineering*, *11*(1), 016002. https://doi.org/ 10.1088/1741-2560/11/1/016002
- Weber, M. J., Messing, S. B., Rao, H., Detre, J. A., & Thompson-Schill, S. L. (2014). Prefrontal transcranial direct current stimulation alters activation and connectivity in cortical and subcortical reward systems: A tDCS-fMRI study. *Human Brain Mapping*, 35(8), 3673–3686. https://doi.org/10.1002/hbm.22429
- Welford, A. (1980). Choice reaction time: Basic concepts. *Reaction Times*, 73–128.
- Wiethoff, S., Hamada, M., & Rothwell, J. C. (2014). Variability in response to transcranial direct current stimulation of the motor cortex. *Brain Stimulation*, 7(3), 468–475. https://doi.org/ 10.1016/j.brs.2014.02.003
- Willis, M. L., Costantino, A. I., Nitsche, M. A., Palermo, R., & Rivolta, D. (2019). Anodal tDCS and high-frequency TRNs targeting the occipitotemporal cortex do not always enhance face perception. *Frontiers in Neuroscience*, 13, 78. https://doi. org/10.3389/fnins.2019.00078
- Yerkes, R. M., & Dodson, J. D. (1908). The relation of strength of stimulus to rapidity of habit-formation. *Journal of Comparative Neurology and Psychology*, 18(5), 459–482. https://doi.org/10. 1002/cne.920180503

How to cite this article: Esposito, M., Ferrari, C., Fracassi, C., Miniussi, C., & Brignani, D. (2022). Responsiveness to left-prefrontal tDCS varies according to arousal levels. *European Journal of Neuroscience*, 1–16. <u>https://doi.org/10.1111/ejn.</u> 15584